

Fermion-boson transmutation and comparison of statistical ensembles in one dimension

K. Schönhammer and V. Meden

Institut für Theoretische Physik der Universität Göttingen, Bunsenstr. 9, D-37073 Göttingen, Germany
(9 August 1995)

The theoretical description of interacting fermions in one spatial dimension is simplified by the fact that the low energy excitations can be described in terms of bosonic degrees of freedom. This fermion-boson transmutation (FBT) which lies at the heart of the Luttinger liquid concept is presented in a way which does not require a knowledge of quantum field theoretical methods. As the basic facts can already be introduced for noninteracting fermions they are mainly discussed. As an application we use the FBT to present exact results for the low temperature thermodynamics and the occupation numbers in the microcanonical and the canonical ensemble. They are compared with the standard grand canonical results.

I. INTRODUCTION

The low temperature thermodynamic properties of simple metals can be qualitatively understood in terms of the simple Sommerfeld model¹, which treats the conduction electrons as noninteracting fermions in a box. Typical results are a specific heat *linear* in temperature and a constant spin susceptibility in qualitative agreement with experiments. Landau's *Fermi liquid* theory² rests on the assumption of *quasi-particles* which are in a one-to-one correspondence to noninteracting fermions. This leads to a linear specific heat and a constant spin susceptibility but involves *renormalized* quantities like the effective mass and the quasi-particle interaction parameters^{2,3}, which are difficult to calculate microscopically. The consistency of the approach was shown using perturbation theory to infinite order and more recently by renormalization group techniques⁴.

The problem of interacting fermions simplifies in one dimension. In a pioneering paper⁵ Tomonaga treated the case of a two-body interaction which is long ranged in real space. He showed that the low energy excitations of the noninteracting as well as the interacting system can be described in terms of *noninteracting bosons*⁶. The important idea to solve the case of interacting fermions was the observation that a long range interaction in real space is short ranged in momentum space and therefore only particles and holes in the vicinity of the Fermi points are involved in the interacting ground state and states with a low excitation energy. To obtain his results, Tomonaga *linearized* the energy dispersion around the two Fermi points $\pm k_F$. Luttinger⁷ later used a model with strictly linear energy dispersions which is closely related to the massless Thirring model⁸. The exact solution for the

Luttinger model was presented by Mattis and Lieb⁹. A very elegant method to calculate correlation functions for the model is the bosonization of the fermion field operator^{10,11}. The exponents of the anomalous power law decay of various correlation functions are determined by the *anomalous dimension*, which can be calculated explicitly for the Tomonaga-Luttinger (TL-) model^{7,9,10}. It was an important observation of Haldane¹¹ that the low energy physics of the exactly solvable TL-model provides the *generic* scenario for one-dimensional fermions with repulsive interactions. Like in the Landau Fermi liquid picture² a few parameters completely determine the low energy physics. Generally they are as difficult to calculate as the Landau parameters. In contrast to the higher dimensional case there are additional exactly solvable models for which Haldane's *Luttinger liquid* scenario can be tested and the parameters determining the anomalous dimension can be calculated using the Bethe-ansatz technique¹².

Another important manifestation of Luttinger liquids is called *spin charge separation*, i.e. for low energy excitations the charge and spin degrees are completely decoupled. This shows up, for example, in the spectral function of the one-particle Green's function^{13,14} which largely determines the photoemission spectrum. Recent high resolution photoemission experiments on quasi one-dimensional organic conductors have been interpreted to show Luttinger liquid behaviour¹⁵.

In the present paper we present the basic ideas behind the Luttinger liquid concept in very simple terms, involving only *basic* quantum mechanics and statistical mechanics. We therefore believe that our approach requires less theoretical knowledge than the field theory approach to FBT published in this journal some time ago¹⁶.

In section II we present the fermion-boson transmutation in terms that should be accessible to anybody who has had an elementary course in solid state physics. Especially *no* use is made of the method of second quantization. The concepts introduced in section II are used in section III and IV to compare the different statistical ensembles for noninteracting fermions in a box^{17,18}. In the grand canonical ensemble the constraint of a fixed particle number which makes the calculation complicated in the canonical ensemble is relaxed and the determination of thermodynamical quantities is straightforward. The microcanonical ensemble is seldom used in this context. Our discussion of the basic ideas of the FBT allows us to present a comparison of the three ensembles. We show

explicitly *how* the results approach each other in the limit of infinite box size with constant density, i.e. the thermodynamic limit. For the calculation of the occupation numbers in the canonical ensemble the numerical effort increases quickly with the system size when the elementary approach of section IV is used. We take this as an opportunity to discuss the fermion-boson transmutation again on a more advanced level using the method of second quantization in section V. The techniques which allow the calculation of the properties of the TL-model of *interacting* fermions^{5,7} are introduced again in the context of noninteracting fermions. The power of this method is shown by presenting a closed expression for the occupation number of noninteracting fermions in the canonical ensemble, which allows to discuss the transition to the Fermi function in the thermodynamic limit.

The paper is written such that readers not familiar with the method of second quantization can understand fermion boson transmutation in its elementary form *and* the comparison of the statistical ensembles by reading sections I to IV only.

II. NONINTERACTING FERMIONS IN A BOX

We consider particles in one dimension in a box of length L with fixed boundary conditions $\phi(0) = \phi(L) = 0$. The single particle eigenfunctions and energies are given by

$$\phi_n(x) = \sqrt{2/L} \sin(k_n x), \quad (1)$$

and

$$\epsilon_n \equiv \epsilon(k_n) = \frac{\hbar^2 k_n^2}{2m}, \quad (2)$$

with $k_n = n\pi/L$ and $n \in \mathbb{N}$. Later we will also consider periodic boundary conditions. Then the spacing of the k -values is doubled and negative *and* positive values occur. In the following the spin of the fermions is neglected for simplicity, i.e. we consider *spinless* fermions as in [5,7]. The ground state of N fermions is given by filling the lowest momentum states up to the Fermi wavevector $k_F = n_F \pi/L$ where $n_F = N$ and the excited states can be classified by giving the *occupation numbers* n_{k_n} , which are zero or one. For fixed particle number N the sum of the occupation numbers has to be equal to N . This constraint makes the evaluation of the *canonical* partition function difficult. In the well known procedure of Darwin and Fowler¹⁸ an integral representation for the Kronecker delta describing the constraint is used. If the corresponding integral is evaluated by the method of steepest descent the result is equivalent to the grand canonical calculation, where only the average particle number is fixed as particle exchange with a bath is assumed. In the following we show that in the *low temperature* regime the canonical partition function can be

expressed as the partition function of a photon gas, i.e. massless bosons.

We first summarize the well known grand canonical results¹⁷. The total energy E_{gc} as a function of the temperature T and the chemical potential μ is given by

$$E_{gc} = \sum_n \frac{\epsilon_n}{e^{\beta[\epsilon_n - \mu]} + 1}, \quad (3)$$

where $\beta = 1/(k_B T)$ and k_B is the Boltzmann constant. The chemical potential μ becomes a function of temperature if the average particle number is assumed to be given by the *fixed* value N

$$N = \sum_n \frac{1}{e^{\beta[\epsilon_n - \mu(T)]} + 1}. \quad (4)$$

In the thermodynamic limit $L \rightarrow \infty, N/L$ fixed, the low temperature result for the energy, the specific heat and other thermodynamic quantities can be explicitly evaluated using the Sommerfeld technique^{1,17}. They are determined by the one-particle density of states per unit length $\rho_0(\epsilon) = 1/\pi \hbar v(\epsilon)$ in the *vicinity* of the Fermi energy $\epsilon_F \equiv \epsilon_{n_F}$, where $\hbar$ times the velocity $v(\epsilon)$ is given by $d\epsilon(k)/dk$ as a function of ϵ . The low temperature specific heat, for example, is *linear* in temperature and determined by $\rho(\epsilon_F)$ ^{1,17}. This is an indication that in order to arrive at this result one is allowed to *simplify* the spectrum of eigenvalues in Eq. (2) in the vicinity of the Fermi energy.

For $n \gg 1$ the eigenvalue spectrum given by Eq. (2) becomes locally *equidistant*

$$\frac{\epsilon_{n+1} - \epsilon_n}{\epsilon_n - \epsilon_{n-1}} = \frac{2n+1}{2n-1} \rightarrow 1. \quad (5)$$




FIG. 1. Specific heat for the quadratic (solid line) and the linearized (dashed line) electron dispersions.

As in Refs. [5] and [7] we therefore *linearize* the spectrum ϵ_n around the Fermi point n_F

$$\epsilon_n^{(l)} = \epsilon_F + \hbar v_F (n - n_F) \pi / L, \quad (6)$$

where $v_F = \hbar k_F / m = \hbar \pi (N/L) / m$ is the Fermi velocity. As we keep the density (N/L) constant while increasing L the Fermi velocity is *independent* of the system size. In the following we consider this as the model for which we want to determine the low temperature thermodynamics. Fig. 1 shows the temperature regime $k_B T \ll \epsilon_F$ for which the specific heats for both models agree. The results were obtained by numerical evaluation of Eqs. (3) and (4) in the thermodynamic limit. The linearization in Eq. (6) is the essential step to obtain the exact solution for the *interacting* TL-model^{5,9}. The constant energy separation of the linearized model is given by $\Delta \equiv \hbar v_F \pi / L$.

For the linearized model also the excitation spectrum of the N -electron system is equidistant

$$E_M = E_0 + M\Delta, \quad (7)$$

where E_0 is the ground state energy of the filled Fermi sea. For large M the eigenvalues are *highly degenerate*. The determination of the degeneracy can be reduced to an old problem in combinatorics¹⁹, if we classify the corresponding states not by fermionic occupation numbers as discussed above, but in terms of the *upward shifts of the occupied levels compared to the Fermi sea*⁶. This is exemplified in Fig. 2 where a special excited state for $M = 20$ is shown. The highest occupied level is shifted by seven energy units, the second and third by four, the fourth by three and the fifth and sixth highest by one unit. All deeper levels are unshifted. The excitation energy in units of Δ is given by $M = 7 + 4 + 4 + 3 + 1 + 1 = 20$. This is one of the possible *partitions*^{19–21} of the number $M = 20$.




FIG. 2. Example of the classification scheme for the excited states.

Each other of the 627 partitions of $M = 20$, written in

lexicographic order, as in our example, uniquely classifies an other possible excited state corresponding to this excitation energy. For small M it is easy to write down all partitions. For $M = 4$ the five possible partitions are: $[4]$; $[3, 1]$; $[2, 2]$; $[2, 1, 1]$; $[1, 1, 1, 1]$. For a general M a partition is defined by a set of M numbers $m_i \in \mathbb{N}_0$ such that

$$M = m_1 + 2m_2 + 3m_3 + \dots + Mm_M = \sum_{l=1}^M lm_l \quad (8)$$

Already more than two hundred years ago Euler¹⁹ proved important theorems about the number of partitions. His use of generating functions can be considered the earliest precursor of the methods of quantum statistical mechanics.

A formally simple expression for the number g_M of partitions of M can be given as a sum over a Kronecker delta using the defining Eq. (8)

$$g_M = \sum_{m_1=0}^{\infty} \dots \sum_{m_M=0}^{\infty} \sum_{m_{M+1}=0}^{\infty} \dots \delta_{M, \sum_{l=1}^{\infty} lm_l}. \quad (9)$$

From Eq. (8) it is obvious that it is unnecessary to include more than M summations in Eq. (9), but the form with the additional infinite number of summations turns out to be useful to calculate the canonical partition function of the linearized model

$$Z_c = \sum_{M=0}^{\infty} \tilde{g}_M e^{-\beta(E_0 + M\Delta)}. \quad (10)$$

Here $\tilde{g}_M$ is the degeneracy corresponding to the excitation energy $M\Delta$. For excitation energies $M\Delta$ smaller than $n_F\Delta$ all partitions of M lead to a possible excited state of the N -particle system and $\tilde{g}_M = g_M$. For $M > n_F$ the degeneracy of excited states is smaller than the number of partitions $\tilde{g}_M < g_M$. For $M = n_F + 1$, e.g. there is no allowed excited N -particle state related to the partition

$$\underbrace{[1, 1, \dots, 1]}_{n_F+1},$$

because in the groundstate there are only n_F fermions that can be shifted upwards. For $M > n_F$ we have to take into account that one-particle states exist only for quantum numbers $n \geq 1$. Two arguments can be given to use g_M as the degeneracy also for $M > n_F$. The one in spirit of Tomonaga's⁵ treatment of the interacting case is to argue that for $n_F\Delta / (k_B T) \gg 1$ the contribution of these highly excited states is *exponentially small* in this large ratio and the replacement of $\tilde{g}_M$ by g_M in Eq. (10) makes no relevant difference. The solution corresponding to Luttinger's⁷ treatment of interacting fermions consists of *adding an infinite Dirac sea* of fermions in one-particle states with $n \leq 0$. Then the replacement of $\tilde{g}_M$ by g_M

is exact, but one has to be careful with the treatment of an infinite number of electrons^{7,9}.

After the replacement $\tilde{g}_M \rightarrow g_M$ the calculation of the partition function Z_c in Eq. (10) is simple if we use Eq. (9)

$$\begin{aligned} Z_c &= e^{-\beta E_0} \sum_{M=0}^{\infty} \sum_{m_1=0}^{\infty} \dots \sum_{m_M=0}^{\infty} \dots \delta_{M, \sum_{l=1}^{\infty} l m_l} \\ &\quad \times e^{-\beta \Delta \sum_{l=1}^{\infty} l m_l} \\ &= e^{-\beta E_0} \prod_{l=1}^{\infty} \left(\sum_{m_l=0}^{\infty} e^{-m_l \beta l \Delta} \right) \\ &= e^{-\beta E_0} \prod_{l=1}^{\infty} \frac{1}{1 - e^{-\beta l \Delta}}. \end{aligned} \quad (11)$$

This is the partition function of a system of massless bosons or harmonic oscillators with energy $l\Delta = \hbar v_F k_l$. The m_l which describe the partitions play the roles of *bosonic quantum numbers*. This is FBT in its most elementary form.

If we differentiate $-\ln Z_c$ with respect to β we obtain the canonical energy E_c

$$E_c(T) = E_0 + \sum_{l=1}^{\infty} \frac{l\Delta}{e^{\beta l \Delta} - 1}. \quad (12)$$

The evaluation of Eq. (12) for finite systems will be discussed in the next section. Here we first want to show that the results for the energy per unit length $e(T) \equiv [E(T) - E_0]/L$ using Eq. (3) and Eq. (12) agree in the thermodynamic limit. In this limit the sums can be replaced by integrals and we obtain from Eq. (12)

$$e_c(T) = \frac{(k_B T)^2}{\pi \hbar v_F} I_-, \quad (13)$$

where the integrals $I_{\pm}$ are defined as

$$I_{\pm} \equiv \int_0^{\infty} \frac{x}{e^x \pm 1} dx. \quad (14)$$

Eq. (13) is the one-dimensional version of the Stefan-Boltzmann law¹⁷ of black body radiation which states that in d dimensions the energy density is proportional to T^{d+1} , or to Debye's theory of the low temperature specific heat due to acoustical phonons¹.

For the calculation in the grand canonical ensemble using Eq. (3) the contribution of the states above ϵ_F equals the part below ϵ_F for the linear dispersion and the chemical potential is temperature *independent*. This yields

$$e_{gc}(T) = \frac{(k_B T)^2}{\pi \hbar v_F} 2I_+. \quad (15)$$

The integrals $I_{\pm}$ can be found in every good table of integrals or textbook on statistical mechanics¹⁷. The fact that the results in Eqs. (13) and (15) agree can be shown

without actually calculating the integrals. Elementary algebra yields

$$I_- - I_+ = 2 \int_0^{\infty} \frac{x}{e^{2x} - 1} dx = I_-/2. \quad (16)$$

and we obtain $I_- = 2I_+$, which shows the equivalence of the canonical and the grand canonical ensemble in the thermodynamic limit. The numerical value is given by¹⁷ $I_- = \zeta(2) = \pi^2/6$ where $\zeta(x)$ is the Riemann zeta function. The specific heat per unit length $c_L \equiv C_L/L$ for both ensembles is therefore in the thermodynamic limit given by

$$c_{\infty} = \frac{\pi}{3} \frac{k_B^2 T}{\hbar v_F}. \quad (17)$$

In the following section the specific heat for both ensembles is discussed for finite systems. For a given density and temperature there is a natural length scale (thermal de Broglie wave length) given by $a(T) \equiv \hbar v_F \pi / (k_B T)$. The system size L is large compared to $a(T)$ if $k_B T$ is large compared to the level spacing Δ .

So far we have not discussed the microcanonical ensemble in which one works with a fixed energy¹⁷ and the entropy as a function of energy is determined by the logarithm of the degeneracy. Therefore we have for the linearized model

$$S_{micro}(E_M) = k_B \ln(g_M), \quad (18)$$

where g_M is the number of partitions. There are recursive methods to determine g_M for small and intermediate values of M and excellent estimates can be given for large M ^{20,22}. We show also in the following section how the microcanonical results approach those of the canonical and the grand canonical ensemble for large L .

III. THERMODYNAMIC PROPERTIES FOR FINITE SYSTEMS

The Stefan-Boltzmann law Eq. (13) for the energy per unit length is only correct in the thermodynamic limit. For finite L the sum in Eq. (12) has to be performed numerically. For $L/a(T) \gg 1$ the Euler summation formula in its most simple form²²

$$\sum_{j=j_0}^{j=j_1} f(j) = \int_{j_0}^{j_1} f(x) dx + [f(j_0) + f(j_1)]/2 \mp \dots, \quad (19)$$

provides the leading finite size correction

$$\frac{c_{L,c}}{c_{\infty}} = 1 - \frac{3}{2\pi^2} \frac{a(T)}{L} + \mathcal{O}\left(\left[\frac{a(T)}{L}\right]^2\right). \quad (20)$$

Fig. 3 shows $c_{L,c}/c_{\infty}$ as a function of $L/a(T)$. As $L/a(T) = k_B T / \Delta$ this plot can be viewed as the specific

heat ratio as a function of system size for fixed temperature or as a function of temperature for a given length of the system. For small arguments the ratio is small due to the well known phenomenon of freezing of the excited states for temperatures small compared to the excitation energies¹⁷. Fig. 3 also shows the corresponding results for the *grand canonical* specific heat following from the numerical evaluation of the result using Eq. (3). For small arguments the exponential suppression is weaker than in the canonical case as the chemical potential lies in the *middle* between the highest occupied and the lowest unoccupied state of the Fermi sea. The use of the Euler summation formula Eq. (19) shows that there is *no* finite size correction of order $a(T)/L$ as in the canonical result Eq. (20) and the asymptotic limit is reached very quickly.




FIG. 3. Ratio of the grand canonical specific heat and the specific heat in the thermodynamic limit c_∞ (solid line) and the canonical specific heat and c_∞ (dashed line) as a function of $L/a(T)$.

As discussed in the preceding section the number of partitions g_M plays a central role for the thermodynamics of the linearized model. In the microcanonical ensemble the number enters explicitly in Eq. (18). The values of the g_M for M up to 500 can be found in a standard handbook²². They increase very quickly with M as can be seen from $g_{500} \approx 2.3 \cdot 10^{21}$, which leads to $S_{micro}(E_0 + 500\Delta)/k_B \approx 49.20$. For values of M of the order fifty or larger already the crudest approximation of Ramanujan^{20,22}

$$(g_M)_R = e^{\pi\sqrt{2M/3}}/(4M\sqrt{3}), \quad (21)$$

provides a good approximation. One obtains, for example $\ln\{g_M(500)\}_R \approx 49.18$. In Fig. 4 we show S_{micro} as a function of M and compare the exact result using the recursion relation^{20,22} with the Ramanujan approximation Eq. (21), which smoothes the number theoretical fluctuations. The inverse microcanonical temperature is defined as

$$\frac{1}{T_{micro}(E_M)} = \left[\frac{\partial S_{micro}(E_M)}{\partial E_M} \right]_L = \frac{k_B}{\Delta} [\ln g_M]'. \quad (22)$$

Due to the discreteness of the energy the derivative in Eq. (22) has to be evaluated as a ratio of differences $f'_M = (f_{M+1} - f_{M-1})/2$. This definition produces smoother results than the definition involving differences of neighbouring numbers. The microcanonical definition of the heat capacity for constant L is

$$C_{L,micro}(E_M) = \left[\left(\frac{\partial T_{micro}(E_M)}{\partial E_M} \right)_L \right]^{-1}. \quad (23)$$




FIG. 4. Microcanonical entropy as a function of the energy calculated with help of the exact degeneracies (circles) and the Ramanujan approximation Eq. (18) (solid line).




FIG. 5. Ratio of the exact microcanonical specific heat and the one calculated with the help of the Ramanujan approximation as a function of $[(E_M - E_0)/\Delta]^{1/2}$.

For $M \gg 1$ an analytical approximation for this quantity can be obtained using the Ramanujan expression Eq. (21), which we denote as $C_{L,R}$

$$C_{L,R}(E_M) = \pi k_B \sqrt{\frac{2(E_M - E_0)}{3\Delta}} + \mathcal{O}\left(\sqrt{\frac{\Delta}{E_M - E_0}}\right). \quad (24)$$

This should be compared with the relation $Lc_\infty = \pi k_B \sqrt{2Le(T)/(3\Delta)}$, where $e(T) = e_c(T) = e_{gc}(T)$, which follows from Eqs. (13) or (15). As in the large energy limit $E - E_0$ in the microcanonical ensemble and $E(T) - E_0$ in the canonical (or grand canonical) ensemble can be identified, as discussed in the next section, $C_{L,R}/L$ agrees with c_∞ in this limit. Fig. 5 shows how the exact microcanonical result Eq. (23) approaches $C_{L,R}$ as a function of $\sqrt{(E_M - E_0)/\Delta}$, which takes the role of $L/a(T)$ in Fig. 3. The ratio $c_{L,micro}/c_{L,R}$ shows quite strong fluctuations for values $\sqrt{(E_M - E_0)/\Delta} < 5$.

As in the thermodynamic limit the entropy per unit length agrees for all three ensembles we can use the *grand canonical entropy* to obtain an estimate for the *number of partitions*. This yields the Ramanujan expression Eq. (21) with $4M\sqrt{3}$ replaced by 1.

IV. OCCUPATION NUMBERS

For variable particle number the method of second quantization in Fock space is the appropriate mathematical framework. The Hamiltonian for noninteracting fermions considered in section II reads

$$H_0 = \sum_m \epsilon_m \hat{n}_m = \sum_m \epsilon_m c_m^\dagger c_m, \quad (25)$$

where the $\hat{n}_m$ are the occupation number operators and $c_m^\dagger$ (c_m) are fermionic creation (annihilation) operators, which obey canonical anticommutation relations

$$[c_m, c_l]_+ = 0, \quad [c_m, c_l^\dagger]_+ = \delta_{m,l}. \quad (26)$$

Explicit use of the method of second quantization is only made in the next section. Here we discuss the calculation of the expectation value of $\hat{n}_m$ in the three ensembles. The calculation in the grand canonical ensemble is standard¹⁷ and for arbitrary system size yields the *Fermi function*

$$\langle \hat{n}_m \rangle_{gc,T} = \frac{1}{e^{\beta(\epsilon_m - \mu)} + 1} \equiv f(\epsilon_m). \quad (27)$$

We next consider the model with the linearized dispersion Eq. (6) in the *microcanonical ensemble*. In section II we have labeled the states of excitation energy $M\Delta$ in terms of the *list of upward shifts* $[\alpha]_M$

$$[\alpha]_M = [l_1, l_2, \dots] \text{ with } l_i \geq l_{i+1} \geq 0, \quad (28)$$

which corresponds to a partition in lexicographic order. The expectation value of the occupation number is given by

$$\langle \hat{n}_m \rangle_{micro,E_M} = \frac{1}{g_M} \sum_{[\alpha]_M} \langle [\alpha]_M | \hat{n}_m | [\alpha]_M \rangle. \quad (29)$$

The expectation value of $\hat{n}_m$ in the state $|\alpha]_M\rangle$ can easily be calculated due to the definition of the list $[\alpha]_M$

$$\langle [\alpha]_M | \hat{n}_m | [\alpha]_M \rangle = 1 \text{ for } m = n_F + l_i - i + 1, \quad (30)$$

and zero otherwise. In order to perform the numerical calculations all partitions of M have to be *explicitly* created, which leads to a drastic increase of computer time for increasing M . The numerical results will be discussed below together with the *canonical* result

$$\langle \hat{n}_m \rangle_{c,T} = \frac{1}{Z_c} \sum_{M=0}^{\infty} g_M \langle \hat{n}_m \rangle_{micro,E_M} e^{-\beta(E_0 + M\Delta)}. \quad (31)$$

The canonical result can therefore be obtained directly from the microcanonical one if the temperature is low enough that the probability distribution

$$w_M(T) = g_M e^{-\beta(E_0 + M\Delta)} / Z_c, \quad (32)$$

has decayed to “zero” for values of M for which the calculation of the microcanonical average is still feasible. Note that while the Fermi function can be obtained from the grand canonical potential by a derivative with respect to ϵ_m ¹⁷ such a procedure is *not* possible for the canonical free energy. In order to calculate the canonical average for higher temperatures the method of the *bosonization of the field operator*^{10,11} can be used which is described in the next section. We therefore only consider results for quite low temperatures in this section.

Fig. 6 shows the average occupation numbers for the three ensembles for $\tau \equiv k_B t / \Delta = 3$. The microcanonical curve is for $M = 10$, which roughly corresponds to the maximum of the probability distribution $w_M(T)$ shown in Fig. 7. The overall agreement of the results is quite good already at this small value of τ . Note that the microcanonical result yields the ground state occupancies for $|m - n_F| > M$, e.g. $\langle \hat{n}_m \rangle_{micro,E_M} = 0$ for $m > M + n_F$, while the canonical and the grand canonical results go to zero continuously.




FIG. 6. Occupation numbers for the three ensembles for $\tau = 3$ respectively $M = 10$.




FIG. 7. Probability distribution of the energy for $\tau \equiv k_B T / \Delta = 3$.

V. BOSONIZATION OF FERMIONIC OPERATORS

This section is more technical than the previous ones and some familiarity with the method of second quantization is required. In section II we have shown that the N fermion states of the model with the linearized dispersion are *highly degenerate* and can be labeled by the list of upward shifts corresponding to partitions, which leads to the *bosonic* canonical partition function. We now show that one can choose *linear combinations* of the degenerate states for a given M , which can be described by bosonic creation and annihilation operators (ladder operators). They play the essential role for the solution of the interacting fermion problem^{5,7,9}. One defines operators bilinear in the fermionic operators

$$b_l^\dagger \equiv \frac{1}{\sqrt{l}} \sum_{n=1}^{\infty} c_{n+l}^\dagger c_n, \quad (33)$$

with $l \geq 1$. This is a sum of operators which all shift a fermion upwards by l energy units. Using the fermionic anticommutation rules it is easy to show that $[b_l^\dagger, b_{l'}^\dagger] = 0$ and

$$[b_l, b_l^\dagger] = \frac{1}{l} \sum_{n=1}^l c_n^\dagger c_n. \quad (34)$$

The operator on the rhs counts the number of fermions in the lowest l states at the bottom of the band divided by l . For $l \ll n_F$ and N fermion states which have *no holes at the bottom of the Fermi sea* one can replace the operator on the rhs of Eq. (34) by the unit operator. One

can make this an exact statement if a Dirac sea is added, i.e. the sum in Eq. (33) runs not from 1 but from $-m_0$ with $m_0 \rightarrow \infty$. An analogous discussion for $l \neq l'$ leads to

$$[b_l, b_{l'}^\dagger] = \delta_{l,l'} \mathbb{1}, \quad (35)$$

i.e. the operators defined in Eq. (33) obey *boson commutation* relations.

With the help of these ladder operators we can create an orthonormal basis set of many fermion states labeled by *bosonic quantum numbers* $\{\tilde{m}_l\}$ with $\tilde{m}_l \in \mathbb{N}_0$, if one treats the Fermi sea $|F(N)\rangle$ as the harmonic oscillator ground state

$$|\{\tilde{m}_l\}\rangle = \prod_l \frac{1}{\sqrt{\tilde{m}_l!}} (b_l^\dagger)^{\tilde{m}_l} |F(N)\rangle. \quad (36)$$

All states with $\sum_l l \tilde{m}_l = M$ provide a new basis of states corresponding to the excitation energy $M\Delta$ of H_0 . They are rather complicated *linear combinations* of the states discussed in section II which could be labeled either by fermionic occupation numbers or by the list $[\alpha]_M$ of the upward shifts. If for $M = 4$ we label the $|\alpha\rangle_M$ by $|i\rangle$ with $i = 1, \dots, 5$, in the order in which the partitions of $M = 4$ were presented in section II, we obtain e.g.

$$\begin{aligned} b_4^\dagger |F\rangle &= \frac{1}{2} (|1\rangle - |2\rangle + |4\rangle - |5\rangle) \\ b_3^\dagger b_1^\dagger |F\rangle &= \frac{1}{\sqrt{3}} (|1\rangle - |3\rangle + |5\rangle) \\ \frac{1}{\sqrt{4!}} (b_1^\dagger)^4 |F\rangle &= \frac{1}{\sqrt{24}} (|1\rangle + 3|2\rangle \\ &\quad + 2|3\rangle + 3|4\rangle + |5\rangle). \end{aligned} \quad (37)$$

As long as we consider only noninteracting fermions both types of basis sets are completely well suited for the description. It turns out that for fermions with a two-body interaction the interacting groundstate has a much simpler form in the *bosonic* basis set Eq. (36).

In Eq. (33) we have expressed the Bose operators $b_l(b_l^\dagger)$ in terms of the fermionic operators. In order to take advantage of the bosonization one has to address the reverse problem: *Can one express fermionic operators completely in terms of the Bose operators?* Here one proceeds in two steps. It is possible to *bosonize* the operator H_0 with the help of the Kronig identity¹¹

$$\sum_{l=1}^{\infty} l b_l^\dagger b_l = \sum_{n=1}^{\infty} n c_n^\dagger c_n - \frac{1}{2} (\hat{N}^2 + \hat{N}), \quad (38)$$

where $\hat{N} \equiv \sum_n c_n^\dagger c_n$ is the fermionic particle number operator. For the proof of this relation the boson commutation relations of the $b_l(b_l^\dagger)$ is not needed. As the lhs of Eq. (38) contains products of *four* fermion operators it might look surprising that one obtains a one-particle operator and only a term involving the particle number

operator on the right hand side. It is left as an exercise to the reader to show that the additional four fermion terms exactly vanish. If we subtract $n_F \hat{N}$ and multiply by $\hbar v_F \pi / L$ we obtain

$$H_0 = \hbar v_F \sum_{l=1}^{\infty} \frac{l\pi}{L} b_l^\dagger b_l + h(\hat{N}), \quad (39)$$

where the function $h(\hat{N})$ is irrelevant in the following.

It is also possible to bosonize a long range two-body interaction if *periodic* boundary conditions are used. Generalizing the procedure described above one linearizes around *both* Fermi points $\pm k_F$. Then the Bose operators corresponding to Eq. (33) are the two contributions to the Fourier transformed density operators. As the two-body interaction is *quadratic in the density* the two-body operator is *bilinear* in the Bose operators. The problem of interacting fermions is therefore reduced to a quadratic form in Bose operators, which is easily diagonalized by a canonical transformation^{5,9}. We do not further discuss interacting fermions here but want to calculate the expectation value of $\hat{n}_m$ in the canonical ensemble for noninteracting electrons. For that purpose we have to express $\hat{n}_m$ in terms of Bose operators. In order to do that we have to proceed similarly as in the *bosonization of the field operator*^{10,11}.

This requires some preliminary steps. Let us consider the following operators *linear* in the boson operators

$$A_+ = \sum_{n=1}^{\infty} \alpha_n b_n^\dagger ; \quad B_- = \sum_{n=1}^{\infty} \beta_n b_n, \quad (40)$$

where α_n and β_n are arbitrary constants. Using the boson commutation relations Eq. (35) one can prove the commutation relations

$$\begin{aligned} [b_m, e^{B_-} e^{A_+}] &= \alpha_m e^{B_-} e^{A_+} \\ [b_m^\dagger, e^{B_-} e^{A_+}] &= -\beta_m e^{B_-} e^{A_+}. \end{aligned} \quad (41)$$

If we can construct an operator S in terms of the fermionic operators which fulfills the commutation relations $[b_m, S] = -\alpha_m S$ and $[b_m^\dagger, S] = \beta_m S$ then $S e^{B_-} e^{A_+}$ commutes with all b_m and $b_m^\dagger$ and can therefore only be a function $G_{\hat{N}}$ of the number operator $\hat{N}$, which commutes with the boson operators²³, i.e.

$$S = G_{\hat{N}} e^{-A_+} e^{-B_-}. \quad (42)$$

In order to bosonize $c_k^\dagger c_n$ we consider the more general operator $c_k^\dagger c_l$ with $k, l \in \mathbb{Z}$, i.e. the infinite Dirac sea is included, and determine its commutation relations with b_m and $b_m^\dagger$. Using the anticommutation relations Eq. (26) one obtains

$$\begin{aligned} [b_m, c_k^\dagger c_l] &= \frac{1}{\sqrt{m}} (c_{k-m}^\dagger c_l - c_k^\dagger c_{l+m}) \\ [b_m^\dagger, c_k^\dagger c_l] &= \frac{1}{\sqrt{m}} (c_{k+m}^\dagger c_l - c_k^\dagger c_{l-m}). \end{aligned} \quad (43)$$

If we now multiply Eq. (43) by $e^{-iku} e^{ilv}$ with real u and v and sum over k and l the commutation relations with the resulting operator are of the form as discussed following Eq. (41)

$$\begin{aligned} [b_m, \psi^\dagger(u) \psi(v)] &= \frac{1}{\sqrt{m}} (e^{-imu} - e^{-imv}) \psi^\dagger(u) \psi(v) \\ [b_m^\dagger, \psi^\dagger(u) \psi(v)] &= \frac{1}{\sqrt{m}} (e^{imu} - e^{imv}) \psi^\dagger(u) \psi(v), \end{aligned} \quad (44)$$

and the auxiliary field operator $\psi(v)$ is defined as

$$\psi(v) = \sum_{l=-\infty}^{\infty} e^{ilv} c_l. \quad (45)$$

It is periodic in v with period 2π . Using the arguments presented following Eq. (41) the bosonized form of $\psi^\dagger(u) \psi(v)$ reads

$$\psi^\dagger(u) \psi(v) = G_{\hat{N}}(u, v) e^{-i(\phi^\dagger(u) - \phi^\dagger(v))} e^{-i(\phi(u) - \phi(v))}, \quad (46)$$

where the operator $\phi(u)$ is given by

$$\phi(u) = -i \sum_{n=1}^{\infty} \frac{e^{inu}}{\sqrt{n}} b_n, \quad (47)$$

and $G_{\hat{N}}(u, v)$ is a so far undetermined function of the particle number operator. As the Fermi sea with N fermions in addition to the Dirac sea is the vacuum state for the bosons, i.e. $b_l |F(N)\rangle = 0$, the expectation value of the product of the exponentials in Eq. (46) in the Fermi sea equals one and yields

$$G_N(u, v) = \langle F(N) | \psi^\dagger(u) \psi(v) | F(N) \rangle, \quad (48)$$

which can be calculated in the fermionic picture using the definition Eq. (45) and $\langle F(N) | c_k^\dagger c_l | F(N) \rangle = \delta_{k,l} \Theta(N - l)$, where $\Theta(x)$ is the step function ($\Theta(0) = 1$)

$$\begin{aligned} G_N(u, v) &= \sum_{l=-\infty}^N e^{il(u-v)} \\ &= e^{iN(u-v)} \sum_{l=-\infty}^0 e^{il(u-v-i0)}. \end{aligned} \quad (49)$$

In the second equality we have added an infinitesimal imaginary shift in $u - v$ in order to make the formal sum convergent. This is a typical example of the problems with the infinite Dirac sea.

After the bosonization of $\psi^\dagger(u) \psi(v)$ is completed, which actually is simpler than the bosonization of a single field operator as no additional operators changing the fermion number have to be introduced¹¹, we can come to our application of calculating the canonical expectation value of $c_m^\dagger c_m$. If we denote the canonical average of an

operator A in a state with N fermions in addition to the Dirac sea by $\langle A \rangle_{N,T}$ we obtain with the Bose function $b(x) \equiv (e^x - 1)^{-1}$

$$\langle \psi^\dagger(u)\psi(v) \rangle_{N,T} = G_N(u, v) \times \exp \left\{ \sum_{n=1}^{\infty} (2 \cos[n(u-v)] - 2) \frac{b(n\Delta\beta)}{n} \right\}. \quad (50)$$

Here we have used the formula $\langle e^A e^B \rangle = e^{\frac{1}{2}\langle A^2 + 2AB + B^2 \rangle}$ for a canonical expectation value for free bosons which was proved by Mermin²⁴ in “one sentence”. It is valid for operators A and B which are *linear* in the boson creation and annihilation operators. To calculate the expectation value of the occupancies we need the inversion formula to Eq. (45)

$$c_l = \frac{1}{2\pi} \int_0^{2\pi} e^{-i v l} \psi(v) dv. \quad (51)$$

As the rhs of Eq. (50) is a function of $u - v$ only one integration can be performed and one obtains

$$\langle c_{N+m}^\dagger c_{N+m} \rangle_{N,T} = \frac{1}{2\pi} \int_0^{2\pi} e^{-i u (m+N)} \times \langle \psi^\dagger(u)\psi(0) \rangle_{N,T}. \quad (52)$$

In order to analytically perform the remaining integration in Eq. (52) we expand $\langle \psi^\dagger(u)\psi(0) \rangle$ in a Laurent series in $z \equiv \exp(iu)$

$$\langle \psi^\dagger(u)\psi(0) \rangle_{N,T} = e^{i u N} \sum_{n=-\infty}^{\infty} d_n e^{i u n}, \quad (53)$$

where the coefficients d_n can be calculated recursively as discussed in the appendix²¹. Using the notation of section IV we finally obtain

$$\langle \hat{n}_{n_F+m} \rangle_{c,T} = d_m. \quad (54)$$

For all values of τ for which the calculation of $\langle \hat{n}_m \rangle_{c,T}$ using Eq. (31) is numerically feasible (e.g. $\tau = 3$ in Fig. 6), the results using Eqs. (31) and (54) are identical. As mentioned in the introduction the result of the present section which used Eq. (46) as the essential ingredient is *very* close in spirit to the calculation of the occupancies of the *interacting* TL-model.




FIG. 8. The difference between the canonical occupation numbers and the grand canonical Fermi function at $\tau = 10$ (solid line) and $\tau = 15$ (dashed line).

Fig. 8 shows the difference of the canonical and the grand canonical result for $\tau = 10$ and 15 as a function of $m - n_F$. From a plot of the maximum deviation as a function of $1/\tau$ we find that it scales as $1/\tau$, i.e. $1/L$ if the temperature T is kept constant. The relative deviation of the canonical occupation numbers from the Fermi function $|\langle n_m \rangle_c - f(\epsilon_m)|/f(\epsilon_m)$ for $m > n_F$ increases with $m - n_F$ and saturates in the exponential tail of the Fermi function. This shows that the canonical occupation numbers decay as $(1 - \eta_\tau) \exp\{-(m - n_F - 0.5)/\tau\}$ for $(m - n_F)/\tau \gg 1$. For $\tau = 15$, for example, $\eta_\tau = 0.032$ and η_τ goes to zero like $1/\tau$. In the thermodynamic limit $L \rightarrow \infty$ the sum in the exponent on the rhs of Eq. (50) goes over to an integral which can be performed analytically. Performing the following integration in Eq. (52) one obtains the Fermi function, i.e. as expected and seen in Fig. 8 the canonical occupation numbers approach the grand canonical ones for $L \rightarrow \infty$.

SUMMARY

This paper intended to serve a dual purpose. First to explain in terms as simple as possible the *concept of bosonization* which plays the essential role in the Luttinger liquid picture of interacting fermions in one dimension. We only shortly discussed interacting electrons, but every reader who followed the algebra of section V should have no problem understanding e.g. the calculation of the occupancies in the interacting case¹¹. As an *application of the concepts* we presented results for the microcanonical and the canonical ensemble for noninteracting fermions. The comparison of our results with the standard textbook grand canonical results can serve as a rare *explicit* example for the general arguments concerning the equivalence of the three ensembles in the thermodynamic limit.

APPENDIX A:

In this appendix we present the method to obtain the coefficients d_m of the Laurent series in Eq. (53) which determine the canonical occupation numbers $\langle n_{n_F+m} \rangle_{c,T}$. According to Eqs. (49) and (50) we can write

$$\langle \psi^\dagger(u) \psi(0) \rangle_{N,T} = e^{iNu} \left\{ \sum_{l=0}^{\infty} z^{-l} \right\} \times \exp \{ f(z) + f(1/z) - 2f(1) \}, \quad (\text{A1})$$

where $z = \exp(iu)$ and the function $f(z)$ is given by

$$f(z) = \sum_{n=1}^{\infty} \frac{b(n/\tau)}{n} z^n. \quad (\text{A2})$$

The geometric series Eq. (A1) is considered as a formal power series. The first step is to determine the coefficients of the power series of the function

$$F(z) \equiv e^{f(z)} = \sum_{m=0}^{\infty} c_m z^m. \quad (\text{A3})$$

Using $F'(z) = f'(z)F(z)$ and $c_0 = 1$ one can obtain recursion relations for $m \geq 1$

$$c_m = \frac{1}{m} \sum_{l=1}^m b(l/\tau) c_{m-l}. \quad (\text{A4})$$

The next step is to write the last factor in Eq. (A1) as a Laurent series

$$F(z)F(1/z)/[F(1)]^2 \equiv \sum_{l=-\infty}^{\infty} a_l z^l. \quad (\text{A5})$$

By comparing coefficients one obtains using Eqs. (A2) and (A3)

$$a_l = \exp \left\{ -2 \sum_{n=0}^{\infty} b(n/\tau)/n \right\} \sum_{m=0}^{\infty} c_m c_{m+l} = a_{-l}. \quad (\text{A6})$$

As for fixed τ the c_m go to zero exponentially for large m the infinite sum converges well. In the last step we have to multiply the Laurent series in Eq. (A5) with the geometric series in Eq. (A1). This yields with Eqs. (53) and (54)

$$d_m = \sum_{n=0}^{\infty} a_{n+m} = \langle n_{n_F+m} \rangle_{c,T}. \quad (\text{A7})$$

The numerical evaluation of the canonical occupation numbers with this method is possible for *much* higher temperatures than via Eq. (31).

- ¹ N. W. Ashcroft and N. D. Mermin, *Solid State Physics* (Holt, Rinehart and Winston, New York, 1976)
- ² L. D. Landau, "The theory of a Fermi liquid," *Sov. Phys. JETP* **3**, 920-925 (1957)
- ³ D. Pines and P. Nozieres, *The Theory of Quantum liquids* (W. A. Benjamin, New York, 1966)
- ⁴ for a review see R. Shankar, "Renormalization-group approach to interacting fermions," *Rev. Mod. Phys.* **66**, 129-192 (1994)
- ⁵ S. Tomonaga, "Remark on Bloch's method of sound waves applied to many fermion problems," *Progr. Theor. Phys.* **5**, 544-569 (1950)
- ⁶ Reprints of most of the important papers in this field together with an excellent introduction are presented in *Bosonization*, Editor M. Stone (World scientific, Singapore, 1994)
- ⁷ J. M. Luttinger, "An exactly soluble model of a many-fermion system," *J. Math. Phys.* **4**, 1154-1162 (1963)
- ⁸ W. Thirring, "A soluble relativistic field theory," *Ann. Phys. (NY)* **3**, 91-112 (1958)
- ⁹ D. C. Mattis and E. H. Lieb, "Exact solution to a many-fermion system and its associated boson field," *J. Math. Phys.* **6**, 304-312 (1965)
- ¹⁰ A. Luther and I. Peschel, "Single particle states, Kohn anomaly and pairing fluctuations in one dimension," *Phys. Rev. B* **9**, 2911-2919 (1974)
- ¹¹ F. D. Haldane, " 'Luttinger liquid theory' of one dimensional quantum fluids: I. Properties of the Luttinger model and their extension to the general 1D interacting spinless Fermi gas," *J. Phys. C* **14**, 2585-2609 (1981)
- ¹² H. J. Schulz, "Correlation exponents and the metal-insulator transition in the one-dimensional Hubbard model," *Phys. Rev. Lett.* **64**, 2831-2834 (1990)
- ¹³ V. Meden and K. Schönhammer, "Spectral functions for the Tomonaga-Luttinger model," *Phys. Rev. B* **46**, 15753-15760 (1992)
- ¹⁴ J. Voit, "Charge-spin separation and spectral properties of Luttinger liquids," *Phys. Rev. B* **47**, 6740-6743 (1993)
- ¹⁵ B. Dardel, D. Malterre, M. Grioni, P. Weibel, Y. Baer, J. Voit, and D. Jérôme, "Possible observation of a Luttinger-liquid behaviour from photoemission spectroscopy of one-dimensional organic conductors," *Europhys. Lett.* **24**, 687-692 (1993)
- ¹⁶ H. Galic, "Alchemy in 1+1 dimension: From bosons to fermions," *Am. J. Phys.* **59**, 1088-1096 (1991)
- ¹⁷ e.g. F. Reif, *Fundamentals of Statistical and Thermal Physics* (MacGraw-Hill, New York, 1965)
- ¹⁸ D. ter Haar, *Elements of Statistical Mechanics* (Rinehart, New York 1954)
- ¹⁹ L. Euler, "De partitione numerorum," (1753) in *Leonardi Euleri opera omnia* (Teubner, Leipzig, 1911)
- ²⁰ G. E. Andrews, *The Theory of Partitions* (Addison-Wesley, Reading, 1976)
- ²¹ Partitions were used in the context of calculating spectral functions in the interacting system in: K. Schönhammer and V. Meden, "Nonuniversal spectral properties of the Luttinger model," *Phys. Rev. B* **47**, 16205-16215 (1993)
- ²² M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions* (Dover, New York, 1970)
- ²³ A similar argument is well known in group theory and is

called “Schur’s Lemma”, see e.g. M. Tinkham, *Group Theory and Quantum Mechanics* (MacGraw-Hill, New York, 1964)

²⁴ N. D. Mermin, “A short simple evaluation of expressions of the Debye-Waller form,” J. Math. Phys. **7**, 1038 (1966)